Avian influenza prevalence in Alaskan shorebirds revisited: Insights from seroprevalence samples





Intercontinental movement



Prevalence data



Avian influenza sampling of target species in Alaska: rRT-PCR



	2006		200	7	200	8	200)9	20′	10	Tot	al
	n	+	n	+	n	+	n	+	n	+	n	+
DUNL	898	2	700	0	927	0	670	0	432	0	3627	2
ROSA	173	0	265	0	272	0	217	0	101	0	1028	0
RUTU	30	0	12	0	13	0	14	0	73	1	142	1
SHAS	225	0	52	0	48	0	54	0	14	0	393	0
Total	1326	2	1029	0	1260	0	955	0	620	1	5190	3

Taken from: USFWS/USGS Progress Reports: Sampling for highly pathogenic Asian H5N1 avian influenza in migratory birds in Alaska (2006-2010 field seasons).



Avian influenza prevalence in shorebirds around the globe (rRT-PCR techniques)

Location	Samples	Prevalence	Season
Alaska: Ip et al.	3,180	0.45%	Spring-Fall
Alaska: Winker et al.	1,050	0.0%	Spring
Pacific Coast: Iverson et al.	2,773	0.5%	Spring-Fall
Pacific Coast: Dusek et al.	2,010	0.0%	Spring-Fall
South America: D'Amico et al.	156	0.0%	Winter
South America: Douglas et al.	128	0.0%	Summer-Fall
South America: Escuerdo et al.	165	0.0%	Fall-Winter
South America: Ghersi et al.	505	0.7%	Summer-Winter




Avian influenza prevalence in shorebirds around the globe, cont'd

Location	Samples Prevalence		Season	
Australia: Haynes et al.	3,748	0.05%	Winter	
Australia: Hurt et al.	173	15%	Winter	
New Zealand: Langstaff et al.	801	0.0%	Fall-Winter	
Europe: Munster et al	2,924	0.0%	Various	
Europe: Hesterberg et al.	748	0.2%	Spring-Fall	
South Korea: Munster et al.	5	20%	?	



Delaware Bay, Eastern US






Average prevalence across 16 years = 14%



Virus genetics:

 Limited evidence for trans-hemispheric virus gene transfer on the East Coast, but lack of positives are hindering a thorough analysis of Pacific Coast

shorebirds.






Future work using seroprevalence techniques?





Why seroprevalence?

- Seroprevalence techniques entail sampling blood for antibodies to AI viruses
- Seroprevalence samples provide evidence of infection spanning a longer time period than rRT-PCR samples ('historical' infection vs. current infection)





Seroprevalence Sampling Methods



- .5 ml blood / bird from jugular
- Paired with oral / cloacal swabs using rRT-PCR analysis
- Pribilof Islands (1-16 Aug 2010), Y-K
 Delta (22 Aug-9 Sept 2010)
- 5 species





Dedicated focus on Ruddy Turnstones: Pribilof Islands



Whoosh net at St. George carcass dump



Special Agent Maftei



Results

Species	Site	n	Serology +
Wandering Tattler	St. George	2	0
Ruddy Turnstone	St. Paul	2	0
	St. George	51	7 (13.7%)
	Y-K Delta	5	0
Sharp-tailed Sandpiper	Y-K Delta	3	0
Rock Sandpiper	St. George	21	0
	Y-K Delta	26	0
Dunlin	Y-K Delta	153	4 (2.6%)
Total		263	11 (4.2%)



Results

Species	Site	n	Serology	n	rRT-PCR
Wandering Tattler	St. George	2	0	2	0
Ruddy Turnstone	St. Paul	2	0	0	0
	St. George	51	7 (13.7%)	51	1 (2%)
	Y-K Delta	5	0	5	0
Sharp-tailed Sandpiper	Y-K Delta	3	0	3	0
Rock Sandpiper	St. George	21	0	21	0
	Y-K Delta	26	0	26	0
Dunlin	Y-K Delta	153	4 (2.6%)	153	0
Total		263	11 (4.2%)	261	1 (0.4%)



Results

- All detections were of LPAI
- 71% (187 birds) of all samples were from AHY birds
- 5 of 7 RUTU and 4 of 4 DUNL seropositives from AHY birds





Comparison with other shorebird seroprevalence information

Species	n	% Seropositive			
Ruddy Turnstone	223	65%			
Red Knot	151	53.6%			
Sanderling	128	3.1%			
Dunlin	17	35.3%			
Short-billed Dowitcher	20	5%			
All samples from Delaware Bay region in May					

From: Brown et al. 2010. Prevalence of antibodies to type A influenza virus in wild avian species using two serological assays. Journal of Wildlife Diseases 46:896-911.



Delaware Bay




Fig. 1. Seasonal distribution of influenza A virus isolations from shorebirds and gulls.

Kawaoka et al. 1988

- What is different about the Pacific Coast?
- Timing off, missing key locations, or is prevalence really that much lower?



Acknowledgements



Shanti Davis, Mark Maftei, Callie Gesmundo, Katie Weber, Tjibbe Stelwagen, and Doug Steiner

